

Chapter 10

Relativity for cyclists



WHAT IF THE LAWS OF MOTION retain their form for a family of coordinate frames related by *continuous* symmetries? The notion of ‘fundamental domain’ is of no use here. If the symmetry is continuous, the dynamical system should be reduced to a lower-dimensional, desymmetrized system, with ‘ignorable’ coordinates eliminated (but not forgotten).

We shall describe here two ways of reducing a continuous symmetry. In the ‘method of slices’ or ‘moving frames’ of sect. 10.4 we slice the state space in such a way that an entire class of symmetry-equivalent points is represented by a single point. In the Hilbert polynomial basis approach of sect. 10.5 we replace the equivariant dynamics by the dynamics rewritten in terms of invariant coordinates. In either approach we retain the option of computing in the original coordinates, and then, when done, projecting the solution onto the symmetry reduced state space.

Instead of writing yet another tome on group theory, in what follows we continue to serve group theoretic nuggets on need-to-know basis, through a series of pedestrian examples (but take a slightly higher, cyclist road in the text proper).

10.1 Continuous symmetries

First of all, why worry about continuous symmetries? Here is an example of the effect a continuous symmetry has on dynamics (for physics background, see remark 10.2). exercise 10.1
exercise 10.2

Example 10.1 Complex Lorenz flow: Consider a complex generalization of Lorenz equations (2.12),

$$\begin{aligned} \dot{x} &= -\sigma x + \sigma y, & \dot{y} &= (\rho - z)x - ay \\ \dot{z} &= (xy^* + x^*y)/2 - bz, \end{aligned} \tag{10.1}$$




Figure 10.1: A typical $\{x_1, x_2, z\}$ trajectory of the complex Lorenz flow, with a short trajectory of figure 10.4 whose initial point is close to the relative equilibrium TW_1 superimposed. See also figure 10.7. (R. Wilczak)

where x, y are complex variables, z is real, while the parameters σ, b are real and $\rho = \rho_1 + i\rho_2, a = 1 - ie$ are complex. Recast in real variables, this is a set of five coupled ODEs

$$\begin{aligned} \dot{x}_1 &= -\sigma x_1 + \sigma y_1 \\ \dot{x}_2 &= -\sigma x_2 + \sigma y_2 \\ \dot{y}_1 &= (\rho_1 - z)x_1 - \rho_2 x_2 - y_1 - e y_2 \\ \dot{y}_2 &= \rho_2 x_1 + (\rho_1 - z)x_2 + e y_1 - y_2 \\ \dot{z} &= -bz + x_1 y_1 + x_2 y_2. \end{aligned} \tag{10.2}$$

In all numerical examples that follow, the parameters will be set to $\rho_1 = 28, \rho_2 = 0, b = 8/3, \sigma = 10, e = 1/10$, unless explicitly stated otherwise. As we shall show in example 10.7, this is a dynamical system with a continuous (but no discrete) symmetry. Figure 10.1 offers a visualization of its typical long-time dynamics. It is a mess. In the rest of this chapter we shall investigate various ways of ‘quotienting’ its $SO(2)$ symmetry, and reducing the dynamics to a 4-dimensional reduced state space. As we shall show here, the dynamics has a nice ‘stretch & fold’ action, but that is totally masked by the continuous symmetry drifts. We shall not rest until we attain the simplicity of figure 10.12, and the bliss of 1-dimensional return map of figure 10.14.

In a presence of a continuous symmetry an orbit can explore the manifold swept by combined action of the dynamics and the symmetry induced drifts. Further problems arise when we try to determine whether an orbit shadows another orbit (see the figure 13.1 for a sketch of a close pass to a periodic orbit), or develop symbolic dynamics (partition the state space, as in chapter 11): here a 1-dimensional trajectory is replaced by a $(N+1)$ -dimensional ‘sausage,’ a dimension for each continuous symmetry (N being the total number of parameters specifying the continuous transformation, and ‘1’ for the time parameter t). How are we to measure distances between such objects? We shall develop here more illuminating visualizations of such flow than figure 10.1, learn how to ‘quotient’ their symmetries, and offer computationally straightforward methods of reducing the dynamics to lower-dimensional, reduced state spaces. The methods should also be applicable to high-dimensional flows, such as translationally invariant fluid flows bounded by pipes or planes (see example 10.4).

But first, a lightning review of the theory of Lie groups. The group-theoretical concepts of sect. 9.1 apply to compact continuous groups as well, and will not be repeated here. All the group theory that we shall need is in principle contained in the *Peter-Weyl theorem*, and its corollaries: A compact Lie group G is completely reducible, its representations are fully reducible, every compact Lie group is a closed subgroup of a unitary group $U(n)$ for some n , and every continuous, unitary, irreducible representation of a compact Lie group is finite dimensional.

Example 10.2 Special orthogonal group $SO(2)$ is a group of length-preserving rotations in a plane. ‘Special’ refers to requirement that $\det g = 1$, in contradistinction to the orthogonal group $O(n)$ which allows for $\det g = \pm 1$. A group element can be parameterized by angle θ , with the group multiplication law $g(\theta')g(\theta) = g(\theta' + \theta)$, and its action on smooth periodic functions $u(\theta + 2\pi) = u(\theta)$ generated by

$$g(\theta') = e^{\theta' \mathbf{T}}, \quad \mathbf{T} = \frac{d}{d\theta}. \tag{10.3}$$

Expand the exponential, apply it to a differentiable function $u(\theta)$, and you will recognize a Taylor series. So $g(\theta')$ shifts the coordinate by θ' , $g(\theta') u(\theta) = u(\theta' + \theta)$.

Example 10.3 Translation group: Differential operator \mathbf{T} in (10.3) is reminiscent of the generator of translations. Indeed, for a constant velocity field $v(x) = c \cdot \mathbf{T}$ the time evolution is nothing but a translation by (time \times velocity):

$$e^{-\tau c \cdot \mathbf{T}} u(x) = e^{-\tau c \cdot \frac{d}{dx}} u(x) = u(x - \tau c). \tag{10.4}$$

As x is a point in the Euclidean \mathbb{R}^d space, the group is not compact. In general, a sequence of time steps in time evolution always forms an Abelian Lie group, albeit never as trivial as this free ballistic motion.

If the group actions consist of N rotations which commute, for example act on an N -dimensional cell with periodic boundary conditions, the group is an Abelian group that acts on a torus T^N .

Example 10.4 Continuous symmetries of the plane Couette flow. (continued from example 9.5) The plane Couette flow is a Navier-Stokes flow bounded by two countermoving planes, in a cell periodic in streamwise and spanwise directions. Every solution of Navier-Stokes equations belongs, by the $SO(2) \times SO(2)$ symmetry, to a 2-torus T^2 of equivalent solutions. Furthermore these tori are interrelated by a discrete D_2 group of spanwise and streamwise flips of the flow cell. (continued in example 10.10)

Let G be a group, and $g\mathcal{M} \rightarrow \mathcal{M}$ a group action on the state space \mathcal{M} . The $[d \times d]$ matrices g acting on vectors in the d -dimensional state space \mathcal{M} form a linear representation of the group G . If the action of every element g of a group G commutes with the flow

$$gv(x) = v(gx), \quad gf^t(x) = f^t(gx), \tag{10.5}$$




Figure 10.2: The group orbit $\mathcal{M}_{x(0)}$ of state space point $x(0)$, and the group orbit $\mathcal{M}_{x(t)}$ reached by the trajectory $x(t)$ time t later. Any point on the manifold $\mathcal{M}_{x(t)}$ is physically equivalent to any other.

G is a symmetry of the dynamics, and, as in (9.6), the dynamics is said to be invariant under G , or G -equivariant.

In order to explore the implications of equivariance for the solutions of dynamical equations, we start by examining the way a compact Lie group acts on state space \mathcal{M} . For any $x \in \mathcal{M}$, the group orbit \mathcal{M}_x of x is the set of all group actions (see page 148 and figure 10.2)

$$\mathcal{M}_x = \{gx \mid g \in G\}. \tag{10.6}$$

As we saw in example 10.3, the time evolution itself is a noncompact 1-parameter Lie group. Thus the time evolution and the continuous symmetries can be considered on the same Lie group footing. For a given state space point x a symmetry group of N continuous transformations together with the evolution in time sweeps out, in general, a smooth $(N+1)$ -dimensional manifold of equivalent solutions (if the solution has a nontrivial symmetry, the manifold may have a dimension less than $N + 1$). For solutions p for which the group orbit of x_p is periodic in time T_p , the group orbit sweeps out a compact invariant manifold \mathcal{M}_p . The simplest example is the $N = 0$, no symmetry case, where the invariant manifold \mathcal{M}_p is the 1-torus traced out by a periodic trajectory p . If \mathcal{M} is a smooth C^∞ manifold, and G is compact and acts smoothly on \mathcal{M} , the reduced state space can be realized as a ‘stratified manifold,’ meaning that each group orbit (a ‘stratum’) is represented by a point in the reduced state space, see sect. 10.4. Generalizing the description of a non-wandering set of sect. 2.1.1, we say that for flows with continuous symmetries the non-wandering set Ω of dynamics (2.2) is the closure of the set of compact invariant manifolds \mathcal{M}_p . Without symmetries, we visualize the non-wandering set as a set of points; in presence of a continuous symmetry, each such ‘point’ is a group orbit.

10.1.1 Lie groups for pedestrians

[...] which is an expression of consecration of ‘angular momentum.’

— Mason A. Porter’s student

Definition: A Lie group is a topological group G such that (i) G has the structure of a smooth differential manifold, and (ii) the composition map $G \times G \rightarrow G : (g, h) \rightarrow gh^{-1}$ is smooth, i.e., C^∞ differentiable.

Do not be mystified by this definition. Mathematicians also have to make a living. Historically, the theory of compact Lie groups that we will deploy here emerged as a generalization of the theory of $SO(2)$ rotations, i.e., Fourier analysis. By a ‘smooth differential manifold’ one means objects like the circle of angles that parameterize continuous rotations in a plane, example 10.2, or the manifold swept by the three Euler angles that parameterize $SO(3)$ rotations.

An element of a compact Lie group continuously connected to identity can be written as

$$g(\theta) = e^{\theta \cdot \mathbf{T}}, \quad \theta \cdot \mathbf{T} = \sum \theta_a \mathbf{T}_a, \quad a = 1, 2, \dots, N, \quad (10.7)$$

where $\theta \cdot \mathbf{T}$ is a *Lie algebra* element, and θ_a are the parameters of the transformation. Repeated indices are summed throughout this chapter, and the dot product refers to a sum over Lie algebra generators. The Euclidian product of two vectors x, y will be indicated by x -transpose times y , i.e., $x^T y = \sum_i^d x_i y_i$. Unitary transformations $\exp(\theta \cdot \mathbf{T})$ are generated by sequences of infinitesimal steps of form

$$g(\delta\theta) \approx 1 + \delta\theta \cdot \mathbf{T}, \quad \delta\theta \in \mathbb{R}^N, \quad |\delta\theta| \ll 1, \quad (10.8)$$

where \mathbf{T}_a , the *generators* of infinitesimal transformations, are a set of linearly independent $[d \times d]$ anti-hermitian matrices, $(\mathbf{T}_a)^\dagger = -\mathbf{T}_a$, acting linearly on the d -dimensional state space \mathcal{M} . In order to streamline the exposition, we postpone discussion of combining continuous coordinate transformations with the discrete ones to sect. 10.2.1.

exercise 10.3

For continuous groups the Lie algebra, i.e., the set of N generators \mathbf{T}_a of infinitesimal transformations, takes the role that the $|G|$ group elements play in the theory of discrete groups. The flow field at the state space point x induced by the action of the group is given by the set of N *tangent fields*

$$t_a(x)_i = (\mathbf{T}_a)_{ij} x_j. \quad (10.9)$$

Any representation of a compact Lie group G is fully reducible, and invariant tensors constructed by contractions of \mathbf{T}_a are useful for identifying irreducible representations. The simplest such invariant is

$$\mathbf{T}^T \cdot \mathbf{T} = \sum_\alpha C_2^{(\alpha)} \mathbf{1}^{(\alpha)}, \quad (10.10)$$

where $C_2^{(\alpha)}$ is the quadratic Casimir for irreducible representation labeled α , and $\mathbf{1}^{(\alpha)}$ is the identity on the α -irreducible subspace, 0 elsewhere. The dot product of two tangent fields is thus a sum weighted by Casimirs,

$$t(x)^T \cdot t(x') = \sum_\alpha C_2^{(\alpha)} x_i \delta_{ij}^{(\alpha)} x'_j. \quad (10.11)$$

Example 10.5 $SO(2)$ irreducible representations: (continued from example 10.2) Expand a smooth periodic function $u(\theta + 2\pi) = u(\theta)$ as a Fourier series

$$u(\theta) = \sum_{m=0}^{\infty} (u_1^{(m)} \cos m\theta + u_2^{(m)} \sin m\theta). \quad (10.12)$$

The matrix representation of the $SO(2)$ action (10.3) on the m th Fourier coefficient pair $(u_1^{(m)}, u_2^{(m)})$ is

$$g^{(m)}(\theta') = \begin{pmatrix} \cos m\theta' & \sin m\theta' \\ -\sin m\theta' & \cos m\theta' \end{pmatrix}, \quad (10.13)$$

with the Lie group generator

$$\mathbf{T}^{(m)} = \begin{pmatrix} 0 & m \\ -m & 0 \end{pmatrix}. \quad (10.14)$$

The $SO(2)$ group tangent (10.9) to state space point $u(\theta)$ on the m th invariant subspace is

$$t^{(m)}(u) = \begin{pmatrix} m u_2^{(m)} \\ -m u_1^{(m)} \end{pmatrix}. \quad (10.15)$$

The L^2 norm of $t(u)$ is weighted by the $SO(2)$ quadratic Casimir (10.10), $C_2^{(m)} = m^2$,

$$\oint \frac{d\theta}{2\pi} (\mathbf{T}u(\theta))^T \mathbf{T}u(2\pi - \theta) = \sum_{m=1}^{\infty} m^2 (u_1^{(m)2} + u_2^{(m)2}), \quad (10.16)$$

and converges only for sufficiently smooth $u(\theta)$. What does that mean? We saw in (10.4) that \mathbf{T} generates translations, and by (10.14) the velocity of the m th Fourier mode is m times higher than for the $m = 1$ component. If $|u^{(m)}|$ does not fall off faster than $1/m$, the action of $SO(2)$ is overwhelmed by the high Fourier modes.

Example 10.6 $SO(2)$ rotations for complex Lorenz equations: Substituting the Lie algebra generator


$$\mathbf{T} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (10.17)$$

acting on a 5-dimensional space (10.2) into (10.7) yields a finite angle $SO(2)$ rotation:

$$g(\theta) = \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 & 0 \\ -\sin \theta & \cos \theta & 0 & 0 & 0 \\ 0 & 0 & \cos \theta & \sin \theta & 0 \\ 0 & 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{10.18}$$

From (10.14) we see that the action of $SO(2)$ on the complex Lorenz equations state space decomposes into $m = 0$ G -invariant subspace (z -axis) and $m = 1$ subspace with multiplicity 2.

The generator \mathbf{T} is indeed anti-hermitian, $\mathbf{T}^\dagger = -\mathbf{T}$, and the group is compact, its elements parametrized by $\theta \bmod 2\pi$. Locally, at $x \in \mathcal{M}$, the infinitesimal action of the group is given by the group tangent field $t(x) = \mathbf{T}x = (x_2, -x_1, y_2, -y_1, 0)$. In other words, the flow induced by the group action is normal to the radial direction in the (x_1, x_2) and (y_1, y_2) planes, while the z -axis is left invariant.



fast track:
sect. 10.2, p. 175

10.1.2 Lie groups for cyclists

Here comes all of the theory of Lie groups in one quick serving. You will live even if you do not digest this section, or, to spell it out; skip this section unless you already know the theory of Lie algebras.



The $[d \times d]$ matrices g acting on vectors in the state space \mathcal{M} form a linear representation of the group G . Tensors transform as

$$h'_{ij}{}^k = g_i{}^i{}' g_j{}^j{}' g^k{}_{k'} h_{i'j'}{}^{k'}. \tag{10.19}$$

A multilinear function $h(\bar{q}, \bar{r}, \dots, s)$ is an invariant function if (and only if) for any transformation $g \in G$ and for any set of vectors q, r, s, \dots it is unchanged by the coordinate transformation

$$h(\overline{gq}, \overline{gr}, \dots, gs) = h(\bar{q}, \bar{r}, \dots, s) = h_{ab\dots} \dots^c q^a r^b \dots s_c. \tag{10.20}$$

Examples of such invariant functions are the length $r(x)^2 = \delta_i^j x^i x_j$ and the volume $V(x, y, z) = \epsilon^{ijk} x_i y_j z_k$. Substitute the infinitesimal form of group action (10.8) into (10.19), keep the linear terms. In the index-notation longhand, the Lie algebra generator acts on each index separately,

$$(\mathbf{T}_a)_i{}^j h_{i'j'}{}^{k\dots} + (\mathbf{T}_a)_j{}^i h_{ij'}{}^{k\dots} - (\mathbf{T}_a)_k{}^l h_{ij\dots}{}^{k'} + \dots = 0. \tag{10.21}$$

Hence the tensor $h_{ij\dots}{}^{k\dots}$ is invariant if $\mathbf{T}_a h = 0$, i.e., the N generators \mathbf{T}_a 'annihilate' it.

As one does not want the symmetry rules to change at every step, the generators $\mathbf{T}_a, a = 1, 2, \dots, N$, are themselves invariant tensors:

$$(\mathbf{T}_a)_i{}^j = g^i{}_{i'} g_j{}^{j'} g_{aa'} (\mathbf{T}_{a'})^{i'}{}_{j'}, \tag{10.22}$$

where $g_{ab} = [e^{-i\theta C}]_{ab}$ is the adjoint $[N \times N]$ matrix representation of $g \in G$. The $[d \times d]$ matrices \mathbf{T}_a are in general non-commuting, and from (10.21) it follows that they close N -element Lie algebra

$$[\mathbf{T}_a, \mathbf{T}_b] = \mathbf{T}_a \mathbf{T}_b - \mathbf{T}_b \mathbf{T}_a = -C_{abc} \mathbf{T}_c, \quad a, b, c = 1, 2, \dots, N,$$

where the fully antisymmetric adjoint representation hermitian generators

$$[C_c]_{ab} = C_{abc} = -C_{bac} = -C_{acb}$$

are the structure constants of the Lie algebra.

As we will not use non-Abelian Lie groups in this chapter, we omit the derivation of the Jacobi relation between C'_{abc} 's, and you can safely ignore all this talk of tensors and Lie algebra commutators as far as the pedestrian applications at hand are concerned.

10.1.3 Equivariance under infinitesimal transformations

A flow $\dot{x} = v(x)$ is G -equivariant (10.5) if

exercise 10.5
exercise 10.6

$$v(x) = g^{-1} v(gx), \quad \text{for all } g \in G. \tag{10.23}$$

For an infinitesimal transformation (10.8) the G -equivariance condition becomes

$$v(x) = (1 - \theta \cdot \mathbf{T}) v(x + \theta \cdot \mathbf{T}x) + \dots = v(x) - \theta \cdot \mathbf{T}v(x) + \frac{dv}{dx} \theta \cdot \mathbf{T}x + \dots.$$

The $v(x)$ cancel, and θ_a are arbitrary. Denote the group flow tangent field at x by $t_a(x)_i = (\mathbf{T}_a)_i{}^j x_j$. Thus the infinitesimal, Lie algebra G -equivariance condition is

$$t_a(v) - A(x) t_a(x) = 0, \tag{10.24}$$

where $A = \partial v / \partial x$ is the stability matrix (4.3). If case you find such learned remarks helpful: the left-hand side of (10.24) is the Lie derivative of the dynamical

flow field v along the direction of the infinitesimal group-rotation induced flow $t_a(x) = \mathbf{T}_a x$,

$$\mathcal{L}_{t_a} v = \left(\mathbf{T}_a - \frac{\partial}{\partial y} (\mathbf{T}_a x) \right) v(y) \Big|_{y=x}. \quad (10.25)$$

The equivariance condition (10.24) states that the two flows, one induced by the dynamical vector field v , and the other by the group tangent field t , commute if their Lie derivatives (or the ‘Lie brackets’ or ‘Poisson brackets’) vanish.

exercise 10.7
exercise 10.8
exercise 10.9

Example 10.7 Equivariance of complex Lorenz flow: That complex Lorenz flow (10.2) is equivariant under $SO(2)$ rotations (10.18) can be checked by substituting the Lie algebra generator (10.17) and the stability matrix (4.3) for complex Lorenz flow (10.2),

$$A = \begin{pmatrix} -\sigma & 0 & \sigma & 0 & 0 \\ 0 & -\sigma & 0 & \sigma & 0 \\ \rho_1 - z & -\rho_2 & -1 & -e & -x_1 \\ \rho_2 & \rho_1 - z & e & -1 & -x_2 \\ y_1 & y_2 & x_1 & x_2 & -b \end{pmatrix}, \quad (10.26)$$

into the equivariance condition (10.24). Considering that $t(v)$ depends on the full set of equations (10.2), and $A(x)$ is only its linearization, this is not an entirely trivial statement. For the parameter values (10.2) the flow is strongly volume contracting (4.47),

$$\partial_t v_i = \sum_{i=1}^5 \lambda_i(x, t) = -b - 2(\sigma + 1) = -24 - 2/3, \quad (10.27)$$

at a coordinate-, ρ - and e -independent constant rate.

Checking equivariance as a Lie algebra condition (10.24) is easier than checking it for global, finite angle rotations (10.23).


10.2 Symmetries of solutions

Let $v(x)$ be the dynamical flow, and f^τ the trajectory or ‘time- τ forward map’ of an initial point x_0 ,

$$\frac{dx}{dt} = v(x), \quad x(\tau) = f^\tau(x_0) = x_0 + \int_0^\tau v(x(\tau')) d\tau'. \quad (10.28)$$

As discussed in sect. 9.2, solutions $x(\tau)$ can be classified by their symmetries. Generic trajectories have no symmetry, but recurrent solutions often do. The simplest solutions are the *equilibria* or *steady* solutions (2.8).

Figure 10.3: (a) A *relative equilibrium orbit* starts out at some point $x(0)$, with the dynamical flow field $v(x) = c \cdot t(x)$ pointing along the group tangent space. For the $SO(2)$ symmetry depicted here, the flow traces out the group orbit of $x(0)$ in time $T = 2\pi/c$. (b) An *equilibrium* lives either in the fixed $\text{Fix}(G)$ subspace (x_3 axis in this sketch), or on a group orbit as the one depicted here, but with zero angular velocity c . In that case the circle (in general, N -torus) depicts a continuous family of fixed equilibria, related only by the group action.



Definition: Equilibrium $x_{EQ} \in \mathcal{M}_{EQ}$ is a fixed, time-invariant solution,

$$\begin{aligned} v(x_{EQ}) &= 0, \\ x(x_{EQ}, \tau) &= x_{EQ} + \int_0^\tau d\tau' v(x(\tau')) = x_{EQ}. \end{aligned} \quad (10.29)$$

An *equilibrium* with full symmetry,

$$g x_{EQ} = x_{EQ} \quad \text{for all } g \in G,$$

lies, by definition, in $\text{Fix}(G)$ subspace (9.11), for example the x_3 axis in figure 10.3 (a). The multiplicity of such solution is one. An equilibrium x_{EQ} with symmetry G_{EQ} smaller than the full group G belongs to a group orbit G/G_{EQ} . If G is finite there are $|G|/|G_{EQ}|$ equilibria in the group orbit, and if G is continuous then the group orbit of x is a continuous family of equilibria of dimension $\dim G - \dim G_{EQ}$. For example, if the angular velocity c in figure 10.3 (b) equals zero, the group orbit consists of a circle of (dynamically static) equivalent equilibria.

exercise 10.10
exercise 10.11

Definition: Relative equilibrium solution $x_{TW}(\tau) \in \mathcal{M}_{TW}$: the dynamical flow field points along the group tangent field, with constant ‘angular’ velocity c , and the trajectory stays on the group orbit, see figure 10.3 (a):

$$\begin{aligned} v(x) &= c \cdot t(x), \quad x \in \mathcal{M}_{TW} \\ x(\tau) &= g(-\tau c) x(0) = e^{-\tau c \mathbf{T}} x(0). \end{aligned} \quad (10.30)$$

A *traveling wave*

$$x(\tau) = g(-c\tau) x_{TW} = x_{TW} - c\tau, \quad c \in \mathbb{R}^d \quad (10.31)$$

is a special type of a relative equilibrium of equivariant evolution equations, where the action is given by translation (10.4), $g(y)x(0) = x(0) + y$. A *rotating wave* is

exercise 10.12
exercise 10.13
exercise 10.14
exercise 10.15
exercise 10.16
exercise 10.17
exercise 10.18




Figure 10.4: $\{x_1, x_2, z\}$ plot of the complex Lorenz flow with initial point close to TW_1 . In figure 10.1 this relative equilibrium is superimposed over the strange attractor. (R. Wilczak)

another special case of relative equilibrium, with the action is given by angular rotation. By equivariance, all points on the group orbit are equivalent, the magnitude of the velocity c is same everywhere along the orbit, so a ‘traveling wave’ moves at a constant speed. For an $N > 1$ trajectory traces out a line within the group orbit. As the c_a components are generically not in rational ratios, the trajectory explores the N -dimensional group orbit (10.6) quasi-periodically. In other words, the group orbit $g(\tau)x(0)$ coincides with the dynamical orbit $x(\tau) \in \mathcal{M}_{TW}$ and is thus flow invariant.

Example 10.8 A relative equilibrium: For complex Lorenz equations and our canonical parameter values of (10.2) a computation yields the relative equilibrium TW_1 with a representative group orbit point

$$(x_1, x_2, y_1, 0, z)_{TW_1} = (8.48492, 0.0771356, 8.48562, 0, 26.9999), \quad (10.32)$$

and angular velocity $c_{TW_1} = 1/11$. This corresponds to period $T_{TW_1} = 2\pi/c \approx 69$, so a simulation has to be run up to time of order of at least 70 for the strange attractor in figure 10.1 to start filling in.

Figure 10.4 shows the complex Lorenz flow with the initial point (10.32) on the relative equilibrium TW_1 . It starts out by drifting in a circle around the z -axis, but as the numerical errors accumulate, the trajectory spirals out.


Calculation of the relative equilibrium stability reveals that it is spiral-out unstable, with the very short period $T_{spiral} = 0.6163$. This is the typical time scale for fast oscillations visible in figure 10.1, with some 100 turns for one circumambulation of the TW_1 orbit. In that time an initial deviation from x_{TW_1} is multiplied by the factor $\Lambda_{radial} \approx 500$. It would be sweet if we could eliminate the drift time scale ≈ 70 and focus just on the fast time scale of ≈ 0.6 . That we will attain by reformulating the dynamics in a reduced state space.

Definition: Periodic orbit. Let x be a periodic point on the periodic orbit p of period T ,

$$f^T(x) = x, \quad x \in \mathcal{M}_p.$$

By equivariance, gx is another periodic point, with the orbits of x and gx either identical or disjoint.

Figure 10.5: A periodic orbit starts out at $x(0)$ with the dynamical v and group tangent t flows pointing in different directions, and returns after time T_p to the initial point $x(0) = x(T_p)$. The group orbit of the temporal orbit of $x(0)$ sweeps out a $(1+N)$ -dimensional torus, a continuous family of equivalent periodic orbits, two of which are sketched here. For $SO(2)$ this is topologically a 2-torus.



If gx lands on the same orbit, g is an element of periodic orbit’s symmetry group G_p . If the symmetry group is the full group G , we are back to (10.30), i.e., the periodic orbit is the group orbit traced out by a relative equilibrium. The other option is that the isotropy group is discrete, the orbit segment $\{x, gx\}$ is pre-periodic (or eventually periodic), $x(0) = g_p x(T_p)$, where T_p is a fraction of the full period, $T_p = T/m$, and thus

$$\begin{aligned} x(0) &= g_p x(T_p), & x &\in \mathcal{M}_p, & g_p &\in G_p \\ x(0) &= g_p^m x(m T_p) = x(T) = x(0). \end{aligned} \quad (10.33)$$

If the periodic solutions are disjoint, as in figure 10.5, their multiplicity (if G is finite, see sect. 9.1), or the dimension of the manifold swept under the group action (if G is continuous) can be determined by applications of $g \in G$. They form a family of conjugate solutions (9.10),

$$\mathcal{M}_{g p} = g \mathcal{M}_p g^{-1}. \quad (10.34)$$

Definition: Relative periodic orbit p is an orbit \mathcal{M}_p in state space \mathcal{M} which exactly recurs

$$x_p(0) = g_p x_p(T_p), \quad x_p(\tau) \in \mathcal{M}_p, \quad (10.35)$$

at a fixed relative period T_p , but shifted by a fixed group action g_p which brings the endpoint $x_p(T_p)$ back into the initial point $x_p(0)$, see figure 10.6. The group action g_p parameters $\theta = (\theta_1, \theta_2, \dots, \theta_N)$ are referred to as ‘phases,’ or ‘shifts.’ In contrast to the pre-periodic (10.33), the phase here are irrational, and the trajectory sweeps out ergodically the group orbit without ever closing into a periodic orbit. For dynamical systems with only continuous (no discrete) symmetries, the parameters $\{\theta_1, \theta_2, \dots, \theta_N\}$ are real numbers, ratios π/θ_j are almost never rational, likelihood of finding a periodic orbit for such system is zero, and such relative periodic orbits are almost never eventually periodic.

Relative periodic orbits are to periodic solutions what relative equilibria (traveling waves) are to equilibria (steady solutions). Equilibria satisfy $f^T(x) - x = 0$ and relative equilibria satisfy $f^T(x) - g(\tau)x = 0$ for any τ . In a co-moving frame,

Figure 10.6: A relative periodic orbit starts out at $x(0)$ with the dynamical v and group tangent t flows pointing in different directions, and returns to the group orbit of $x(0)$ after time T_p at $x(T_p) = g_p x(0)$, a rotation of the initial point by g_p . For flows with continuous symmetry a generic relative periodic orbit (not pre-periodic to a periodic orbit) fills out ergodically what is topologically a torus, as in figure 10.5; if you are able to draw such a thing, kindly send us the figure. As illustrated by figure 10.8 (a) this might be a project for Lucas Films.






Figure 10.7: (Figure 10.1 continued) A group portrait of the complex Lorenz equations state space dynamics. Plotted are relative equilibrium TW_1 (red), its unstable manifold (brown), equilibrium EQ_0 , one trajectory from the group orbit of its unstable manifold (green), 3 repetitions of relative periodic orbit \overline{OI} (magenta) and a generic orbit (blue). (E. Siminos)

i.e., frame moving along the group orbit with velocity $v(x) = c \cdot t(x)$, the relative equilibrium appears as an equilibrium. Similarly, a relative periodic orbit is periodic in its mean velocity $c_p = \theta_p/T_p$ co-moving frame (see figure 10.8), but in the stationary frame its trajectory is quasiperiodic. A co-moving frame is helpful in visualizing a single ‘relative’ orbit, but useless for viewing collections of orbits, as each one drifts with its own angular velocity. Visualization of all relative periodic orbits as periodic orbits we attain only by global symmetry reductions, to be undertaken in sect. 10.4.

Example 10.9 Complex Lorenz flow with relative periodic orbit: Figure 10.7 is a group portrait of the complex Lorenz equations state space dynamics, with several important players posing against a generic orbit in the background.

The unstable manifold of relative equilibrium TW_1 is characterized by a 2-dimensional complex eigenvector pair, so its group orbit is a 3-dimensional. Only one representative trajectory on it is plotted in the figure. The unstable manifold of equilibrium EQ_0 has one expanding eigenvalue, but its group orbit is a cone originating at EQ_0 . Only one representative trajectory on this cone is shown in the figure. It lands close to TW_1 , and then spirals out along its unstable manifold. 3 repetitions of a short relative periodic orbit \overline{OI} are drawn. The trajectory fills out ergodically a 2-dimensional orbit M_{01} . The assignment of its symbolic dynamics label will be possible only after the symmetry reduction, see figure 10.14 and figure 11.9.

Figure 10.8: A relative periodic orbit of Kuramoto-Sivashinsky flow projected on (a) the stationary state space coordinate frame $\{v_1, v_2, v_3\}$, traced for four periods T_p ; (b) the co-moving $\{\tilde{v}_1, \tilde{v}_2, \tilde{v}_3\}$ coordinate frame, moving with the mean angular velocity $c_p = \theta_p/T_p$. (from ref. [10.1])



10.2.1 Discrete and continuous symmetries together

We expect to see relative periodic orbits because a trajectory that starts on and returns to a given torus of a symmetry equivalent solutions is unlikely to intersect it at the initial point, unless forced to do so by a discrete symmetry. This we will make explicit in sect. 10.4, where relative periodic orbits will be viewed as periodic orbits of the reduced dynamics.

If, in addition to a continuous symmetry, one has a discrete symmetry which is not its subgroup, one does expect equilibria and periodic orbits. However, a relative periodic orbit can be pre-periodic if it is equivariant under a discrete symmetry, as in (10.33): If $g^m = 1$ is of finite order m , then the corresponding orbit is periodic with period mT_p . If g is not of a finite order, a relative periodic orbit is periodic only after a shift by g_p , as in (10.35). Morally, as it will be shown in chapter 21, such orbit is the true ‘prime’ orbit, i.e., the shortest segment that under action of G tiles the entire invariant submanifold M_p .

Example 10.10 Relative orbits in the plane Couette flow. (continued from example 10.4) Translational symmetry allows for relative equilibria (traveling waves), characterized by a fixed profile Eulerian velocity $u_{TW}(x)$ moving with constant velocity c , i.e.

$$u(x, \tau) = u_{TW}(x - c\tau). \tag{10.36}$$

As the plane Couette flow is bounded by two counter-moving planes, it is easy to see where the relative equilibrium (traveling wave) solutions come from. A relative equilibrium solution hugs close to one of the walls and drifts with it with constant velocity, slower than the wall, while maintaining its shape. A relative periodic solution is a solution that recurs at time T_p with exactly the same disposition of the Eulerian velocity fields over the entire cell, but shifted by a 2-dimensional (streamwise, spanwise) translation g_p . By discrete symmetries these solutions come in counter-traveling pairs $u_q(x - c\tau)$, $-u_q(-x + c\tau)$: for example, for each one drifting along with the upper wall, there is a counter-moving one drifting along with the lower wall. Discrete symmetries also imply existence of strictly stationary solutions, or ‘standing waves.’ For example, a solution with velocity fields antisymmetric under reflection through the midplane has equal flow velocities in opposite directions, and is thus an equilibrium stationary in time.

10.3 Stability

A spatial derivative of the equivariance condition (10.5) yields the matrix equivariance condition satisfied by the stability matrix (stated here both for the finite group actions, and for the infinitesimal, Lie algebra generators):

$$gA(x)g^{-1} = A(gx), \quad [\mathbf{T}_a, A] = \frac{\partial A}{\partial x} t_a(x). \quad (10.37)$$

For a flow within the fixed $\text{Fix}(G)$ subspace, $t(x)$ vanishes, and the symmetry imposes strong conditions on the perturbations out of the $\text{Fix}(G)$ subspace. As in this subspace stability matrix A commutes with the Lie algebra generators \mathbf{T} , the spectrum of its eigenvalues and eigenvectors is decomposed into irreducible representations of the symmetry group. This we have already observed for the EQ_0 of the Lorenz flow in example 9.10.

An infinitesimal symmetry group transformation maps the initial and the end point of a finite trajectory into a nearby, slightly rotated equivalent points, so we expect the perturbations along to group orbit to be marginal, with unit eigenvalues. The argument is akin to (4.7), the proof of marginality of perturbations along a periodic orbit. Consider two nearby initial points separated by an N -dimensional infinitesimal group transformation (10.8): $\delta x_0 = g(\delta\theta)x_0 - x_0 = \delta\theta \cdot \mathbf{T}x_0 = \delta\theta \cdot t(x_0)$. By the commutativity of the group with the flow, $g(\delta\theta)f^T(x_0) = f^T(g(\delta\theta)x_0)$. Expanding both sides, keeping the leading term in $\delta\theta$, and using the definition of the Jacobian matrix (4.6), we observe that $J^T(x_0)$ transports the N -dimensional group tangent space at $x(0)$ to the rotated tangent space at $x(\tau)$ at time τ :

$$t_a(\tau) = J^T(x_0)t_a(0), \quad t_a(\tau) = \mathbf{T}_a x(\tau). \quad (10.38)$$

For a relative periodic orbit, $g_p x(T_p) = x(0)$, at any point along cycle p the group tangent vector $t_a(\tau)$ is an eigenvector of the Jacobian matrix with an eigenvalue of unit magnitude,

$$J_p t_a(x) = t_a(x), \quad J_p(x) = g_p J^{T_p}(x), \quad x \in \mathcal{M}_p. \quad (10.39)$$

Two successive points along the cycle separated by $\delta x_0 = \delta\theta \cdot t(\tau)$ have the same separation after a completed period $\delta x(T_p) = g_p \delta x_0$, hence eigenvalue of magnitude 1. In presence of an N -dimensional Lie symmetry group, N eigenvalues equal unity.

10.4 Reduced state space

Given Lie group G acting smoothly on a C^∞ manifold \mathcal{M} , the *reduced state space* \mathcal{M}/G is the set of equivalence classes, each class a group orbit. In the literature



exercise 10.19
exercise 10.20




Figure 10.9: Two trajectories $x(t), y(t)$ are equivalent up to a group rotation $g(t)$ as long as they belong to the same group orbit $\mathcal{M}_{x(t)}$.

this space is often rediscovered, and thus has many names - it is alternatively called ‘desymmetrized state space,’ ‘orbit space,’ ‘quotient space,’ or ‘image space,’ obtained by mapping equivariant dynamics to invariant dynamics by methods such as ‘moving frames,’ ‘cross sections,’ ‘slices,’ ‘freezing,’ ‘Hilbert bases,’ ‘quotienting,’ or ‘desymmetrization’

remark 10.1

Symmetry reduction replaces a dynamical system (\mathcal{M}, f) with a symmetry by a ‘desymmetrized’ system $(\tilde{\mathcal{M}}, \tilde{f})$, a system where each group orbit is replaced by a point, and the action of the group is trivial, $gy = y$ for all $y \in \tilde{\mathcal{M}}, g \in G$. The reduced state space $\tilde{\mathcal{M}}$ is sometimes called the ‘quotient space’ \mathcal{M}/G because the symmetry has been ‘divided out.’ For a discrete symmetry, the reduced state space \mathcal{M}/G is given by the fundamental domain of sect. 9.4. In presence of a continuous symmetry, the reduction to \mathcal{M}/G amounts to a change of coordinates where the ‘ignorable angles’ $\{\theta_1, \dots, \theta_N\}$ that parameterize N group translations can be separated out.

We start our discussion of symmetry reduction by considering the finite-rotations *method of moving frames*, and its differential formulation, the *method of slices*.

10.4.1 Go with the flow: method of moving frames

The idea: As the symmetries commute with dynamics, we can evolve a solution $x(\tau)$ for as long as we like, and then rotate it to any equivalent point (see figure 10.9) on its group orbit,

$$x(\tau) = g(\tau)y(\tau), \quad (10.40)$$

any time and any way we like. In the ‘method of slices’ one point on each group orbit is picked as the reduced state space representative of the equivalence class, by slicing across the group orbits by a fixed hypersurface. We start by describing how the method works for a finite segment of the full state space trajectory.

Split up the integration into a sequence of finite time steps, each followed by a rigid coordinate frame rotation (‘moving frame’) such that the next segment’s initial point is in the *slice* fixed by a point y' .

Figure 10.10: Slice \tilde{M} is a Poincaré section (10.41) for group orbits (indicated by dotted lines here). The full state space trajectory $x(t)$ and the reduced state space trajectory $y(t)$ belong to the same group orbit $\mathcal{M}_{x(t)}$ and are equivalent up to a group rotation $g(t)$ (10.40).





Figure 10.11: Method of moving frames for a flow $SO(2)$ -equivariant under (10.18) with slice through $y' = (0, 1, 0, 0, 0)$, group tangent $t' = (1, 0, 0, 0, 0)$. The clockwise orientation condition restricts the slice to half-hyperplane $y_1 = 0, y_2 > 0$. A trajectory started on the slice at $y(0)$, evolves to a state space point with a non-zero $x_1(t_1)$. Compute the polar angle θ_1 of $x(t_1)$ in the (x_1, x_2) plane. The entire state space is then rotated (hence ‘moving frame’) clockwise by θ_1 to $y(t_1) = g(-\theta_1)x(t_1)$, so that the equivalent point on the circle lies on the slice, $y_1(t_1) = 0$. Thus after every finite time step followed by a rotation the trajectory restarts from the $y_1(t_i) = 0$ reduced state space.



Definition: Slice. Let G act regularly on a d -dimensional manifold \mathcal{M} , i.e., with all group orbits N -dimensional. A *slice* through point y' is a $(d-N)$ -dimensional submanifold \tilde{M} such that all group orbits in an open neighborhood of the slice-defining point y' intersect \tilde{M} transversally and only once (see figure 10.10).

The simplest *slice condition* defines a slice as a $(d-N)$ -dimensional hyperplane \tilde{M} normal to the N group rotation tangents t'_a at point y' :

$$(y - y')^T t'_a = 0, \quad t'_a = t_a(y') = \mathbf{T}_a y'. \tag{10.41}$$

In other words, ‘slice’ is a Poincaré section (3.6) for group orbits. The G -invariant subspaces are always within the slice, as $\mathbf{T}_a x = 0$ for x in an invariant subspace, see (10.21). Each ‘big circle’ –group orbit tangent to t'_a – intersects the hyperplane exactly twice, with the two solutions separated by π . As for a Poincaré section (3.4), we add an orientation condition, and select the intersection with the clockwise rotation angle into the slice.

Definition: Moving frame. Assume that for a given $x \in \mathcal{M}$ and a given slice \tilde{M} there exists a unique group element $g = g(x)$ that rotates x into the slice, $gx = y \in \tilde{M}$. The map that associates to a state space point x a Lie group action $g(x)$ is called a *moving frame*.

As $y'^T t'_a = 0$ by the antisymmetry of \mathbf{T}_a , the slice condition (10.41) fixes θ for a given x by

$$0 = y'^T t'_a = x^T g(\theta)^T t'_a, \tag{10.42}$$

exercise 6.1
exercise 10.21

where g^T denotes the transpose of g . The method of moving frames is a post-processing method; trajectories are computed in the full state space, then rotated into the slice whenever desired, with the slice condition easily implemented. The slice group tangent t' is a given vector, and $g(\theta)x$ is another vector, linear in x and a function of group parameters θ . Rotation parameters θ are determined numerically, by a Newton method, through the slice condition (10.42).

Figure 10.11 illustrates the method of moving frames for an $SO(2)$ slice normal to the x_2 axis. Looks innocent, except there is nothing to prevent a trajectory from going thorough $(x_1, x_2) = (0, 0)$, and what θ is one to use then? We can always chose a finite time step that hops over this singularity, but in the continuous time formulation we will not be so lucky.

How does one pick a slice point y' ? A generic point y' not in an invariant subspace (on the complex Lorenz equations z axis, for example) should suffice to fix a slice. The rules of thumb are much like the ones for picking Poincaré sections, sect. 3.1.1. The intuitive idea is perhaps best visualized in the context of fluid flows. Suppose the flow exhibits an unstable coherent structure that – approximately – recurs often at different spatial dispositions. One can fit a ‘template’ to one recurrence of such structure, and describe other recurrences as its translations. A well chosen slice point belongs to such dynamically important equivalence class (i.e. group orbit). A slice is locally isomorphic to \mathcal{M}/G , in an open neighborhood of y' . As is the case for the dynamical Poincaré sections, in general a single slice does not suffice to reduce $\mathcal{M} \rightarrow \mathcal{M}/G$ globally.

The Euclidian product of two vectors x, y is indicated in (10.41) by x -transpose times y , i.e., $x^T y = \sum_i^d x_i y_i$. More general bilinear norms $\langle x, y \rangle$ can be used, as long as they are G -invariant, i.e., constant on each irreducible subspace. An example is the quadratic Casimir (10.11).

Note: method of moving frames has nothing in common with the co-moving frames, such as the one illustrated in figure 10.8. Each relative periodic orbit has its own co-moving frame. In the method of moving frames (or the method of slices) one fixes a stationary slice, and rotates all solutions back into the slice.

Example 10.11 An $SO(2)$ moving frame: (continued from example 10.2) The $SO(2)$ action $(y_1, y_2) = (x_1 \cos \theta - x_2 \sin \theta, x_1 \sin \theta + x_2 \cos \theta)$ is regular on $\mathbb{R}^2 \setminus \{0\}$. Thus we can define a slice as a ‘hyperplane’ (here a mere line), through $y' = (0, 1)$, with group tangent $t' = (0, 1)$, and ensure uniqueness by clockwise rotation into positive x_2 axis. Hence the reduced state space is the half-line $x_1 = 0, y_2 = x_2 > 0$. The slice condition $y_1 = 0$ yields the explicit formula for the moving frame parameter

$$\theta(x_1, x_2) = \tan^{-1}(x_1/x_2), \tag{10.43}$$

i.e., the angle which rotates the point (x_1, x_2) back to the slice, taking care that \tan^{-1} distinguishes (x_1, x_2) plane quadrants correctly. As $SO(2)$ preserves lengths, $x_1^2 + x_2^2 = y_1^2 + y_2^2$, and the group-orbit point in the slice is an $SO(2)$ -invariant,

$$y_2 = \sqrt{x_1^2 + x_2^2}. \tag{10.44}$$

The example is a bit trivial. If dynamics is in plane and $SO(2)$ equivariant, the solutions can only be circles of radius $\sqrt{x_1^2 + x_2^2}$, so this is the “rectification” of the harmonic oscillator by a change to polar coordinates, example 6.1. Still, it illustrates the sense in which the method of moving frames yields group invariants. (E. Siminos)

10.4.2 Dynamics within a slice

I made a wrong mistake.
—Yogi Berra

The choice of finite time steps in the preceding sections was arbitrary, so it is tempting to see what happens if the steps are taken infinitesimal. As we shall see, we do get a flow restricted to the slice, but at a price.

By equivariance one can always write the full state space trajectory as $x(\tau) = g(\tau)y(\tau)$, where the $(d-N)$ -dimensional reduced state space trajectory $y(\tau)$ is to be fixed by some condition, and $g(\tau)$ is then the corresponding curve on the N -dimensional group manifold of the group action that rotates y into x at time τ . The time derivative is then $\dot{x} = v(gy) = \dot{g}y + gu$, with the reduced state space velocity field given by $u = dy/dt$. Rewriting this as $u = g^{-1}v(gy) - g^{-1}\dot{g}y$ and using the equivariance condition (10.23) leads to

$$u = v - g^{-1}\dot{g}y.$$

The Lie group element (10.7) and its time derivative describe the group tangent flow

$$g^{-1}\dot{g} = g^{-1}\frac{d}{dt}e^{\theta \cdot \mathbf{T}} = \dot{\theta} \cdot \mathbf{T}.$$

This is the group tangent velocity $g^{-1}\dot{g}y = \dot{\theta} \cdot t(y)$ evaluated at the point y , i.e., with $g = 1$. The flow in the $(d-N)$ directions transverse to the group flow is now obtained by subtracting the flow along the group tangent direction,

$$u(y) = v(y) - \dot{\theta}(y) \cdot t(y), \quad u = dy/dt, \quad (10.45)$$

for any factorization of the flow of form $x(\tau) = g(\tau)y(\tau)$. To integrate these equations we first have to fix a particular flow factorization by imposing conditions on $y(\tau)$, and then integrate phases $\theta(\tau)$ on a given reduced state space trajectory $y(\tau)$.

exercise 10.23
exercise 10.24

Here we demand that the reduced state space is confined to a hyperplane slice. Substituting (10.45) into the time derivative of the fixed slice condition (10.42),

$$u(y)^T t'_a = v(y)^T t'_a - \dot{\theta}_a \cdot t(y)^T t'_a = 0,$$




Figure 10.12: Method of moving frames, slice fixed by a point on the complex Lorenz equations relative equilibrium group orbit, $y' = x_{TW_1}$. (a) The strange attractor of figure 10.1 in the reduced state space of (10.47), $\{x_1, x_2, z\}$ projection. (b) $\{x_2, y_2, z\}$ projection. The reduced state space complex Lorenz flow strange attractor of figure 10.1 now exhibits a discontinuity due to the vanishing denominator in (10.48).

yields the equation for the group phases flow $\dot{\theta}$ for the slice fixed by y' , together with the reduced state space \bar{M} flow $u(y)$:

$$\dot{\theta}_a(y) = \frac{v(y)^T t'_a}{t(y)^T \cdot t'} \quad (10.46)$$

$$u(y) = v(y) - \dot{\theta}(y) \cdot t(y), \quad y \in \bar{M}. \quad (10.47)$$

Each group orbit $\mathcal{M}_x = \{g \cdot x | g \in G\}$ is an equivalence class; method of slices represents the class by its single slice intersection point y . By construction $u^T t' = 0$, and the motion stays in the $(d-N)$ -dimensional slice. We have thus replaced the original dynamical system $\{\mathcal{M}, f\}$ by a reduced system $\{\bar{M}, \bar{f}\}$.

In the pattern recognition and ‘template fitting’ settings (10.46) is called the ‘reconstruction equation.’ Integrated together, the reduced state space trajectory (10.47) and $g(\tau) = \exp\{\theta(\tau) \cdot \mathbf{T}\}$, the integrated phase (10.46), reconstruct the full state space trajectory $x(\tau) = g(\tau)y(\tau)$ from the reduced state space trajectory $y(\tau)$, so no information about the flow is lost in the process of symmetry reduction. exercise 10.25
exercise 10.26

Example 10.12 A slice for complex Lorenz flow. (continuation of example 10.6) Here we can use the fact that

$$t(y)^T \cdot t' = \bar{x}^T \mathbf{T}^T \cdot \mathbf{T} y' = \bar{x}_1 x'_1 + \bar{x}_2 x'_2 + \bar{y}_1 y'_1 + \bar{y}_2 y'_2$$

is the dot-product restricted to the $m = 1$ 4-dimensional representation of $SO(2)$. A generic y' can be brought to form $y' = (0, 1, y'_1, y'_2, z)$ by a rotation and rescaling. Then $\mathbf{T}y' = (1, 0, y'_2, -y'_1, 0)$, and

$$\frac{v(\bar{x}) \cdot t'}{t(\bar{x}) \cdot t'} = \frac{v_1 + v_3 y'_2 - v_4 y'_1}{\bar{x}_2 + \bar{y}_1 y'_1 + \bar{y}_2 y'_2}. \quad (10.48)$$

A long time trajectory of (10.47) with y' on the relative equilibrium TW_1 group orbit is shown in figure 10.12. As initial condition we chose the initial point (10.32) on the unstable manifold of TW_1 , rotated back to the slice by angle θ as prescribed by (10.42). We show the part of the trajectory for $t \in [70, 100]$. The relative equilibrium TW_1 , now an equilibrium of the reduced state space dynamics, organizes the flow into a Rössler type attractor (see figure 2.6). The denominator in (10.46) vanishes and the phase velocity $\dot{\theta}(y)$ diverges whenever the direction of group action on the reduced state space point

is perpendicular to the direction of group action on the slicepoint y' . While the reduced state space flow appears continuous in the $\{x_1, x_2, z\}$ projection, figure 10.12(a), this singularity manifests itself as a discontinuity in the $\{x_2, y_2, z\}$ projection, figure 10.12(b). The reduced state space complex Lorenz flow strange attractor of figure 10.1 now exhibits a discontinuity whenever the trajectory crosses this $-$ -dimensional subspace.

Slice flow equations (10.47) and (10.46) are pretty, but there is a trouble in the paradise. The slice flow encounters singularities in subsets of state space, with phase velocity $\dot{\theta}$ divergent whenever the denominator in (10.48) changes sign, see $\{x_2, y_2, z\}$ projection of figure 10.12(b). Hence a single slice does not in general suffice to cover M/G globally.

10.5 Method of images: Hilbert bases

(E. Siminos and P. Cvitanović)

Erudite reader might wonder: why all this slicing and dicing, when the problem of symmetry reduction had been solved by Hilbert and Weyl nearly a century ago? Indeed, the most common approach to symmetry reduction is by means of a Hilbert invariant polynomial bases (9.15), motivated intuitively by existence of such nonlinear invariants as the rotationally-invariant length $r^2 = x_1^2 + x_2^2 + \dots + x_d^2$, or, in Hamiltonian dynamics, the energy function. One trades in the equivariant state space coordinates $\{x_1, x_2, \dots, x_d\}$ for a non-unique set of $m \geq d$ polynomials $\{u_1, u_2, \dots, u_m\}$ invariant under the action of the symmetry group. These polynomials are linearly independent, but functionally dependent through $m - d + N$ relations called *syzygies*.

Example 10.13 An $SO(2)$ Hilbert basis: (continued from example 9.6) The Hilbert basis

$$\begin{aligned} u_1 &= x_1^2 + x_2^2, & u_2 &= y_1^2 + y_2^2, \\ u_3 &= x_1 y_2 - x_2 y_1, & u_4 &= x_1 y_1 + x_2 y_2, \\ u_5 &= z. \end{aligned} \tag{10.49}$$

is invariant under the $SO(2)$ action on a 5-dimensional state space (10.18). That implies, in particular, that the image of the full state space relative equilibrium TW_{11} group orbit of figure 10.4 is the stationary equilibrium point EQ_1 , see figure 10.13. The polynomials are linearly independent, but related through one syzygy,

$$u_1 u_2 - u_3^2 - u_4^2 = 0, \tag{10.50}$$

yielding a 4-dimensional $M/SO(2)$ reduced state space, a symmetry-invariant representation of the 5-dimensional $SO(2)$ equivariant dynamics. (continued in example 10.14)

The dynamical equations follow from the chain rule

$$\dot{u}_i = \frac{\partial u_i}{\partial x_j} \dot{x}_j, \tag{10.51}$$




Figure 10.13: Invariant ‘image’ of complex Lorenz equations dynamics, figure 10.1, projected onto the invariant polynomial basis (10.49). Note the unstable manifold connection from the equilibrium EQ_0 at the origin to the strange attractor controlled by the rotation around relative equilibrium EQ_1 (the reduced state space image of TW_1); as in the Lorenz flow figure 3.7, natural measure close to EQ_0 is vanishingly small but non-zero.

upon substitution $\{x_1, x_2, \dots, x_d\} \rightarrow \{u_1, u_2, \dots, u_m\}$. One can either rewrite the dynamics in this basis or plot the ‘image’ of solutions computed in the original, equivariant basis in terms of these invariant polynomials.

Example 10.14 Complex Lorenz equations in a Hilbert basis: (continuation of example 10.13) Substitution of (10.2) and (10.49) into (10.51) yields complex Lorenz equations in terms of invariant polynomials:

$$\begin{aligned} \dot{u}_1 &= 2\sigma(u_3 - u_1) \\ \dot{u}_2 &= -2u_2 - 2u_3(u_5 - \rho_1) \\ \dot{u}_3 &= \sigma u_2 - (\sigma - 1)u_3 - e u_4 + u_1(\rho_1 - u_5) \\ \dot{u}_4 &= e u_3 - (\sigma + 1)u_4 \\ \dot{u}_5 &= u_3 - b u_5. \end{aligned} \tag{10.52}$$

As far as visualization goes, we need neither construct nor integrate the invariant dynamics (10.52). It suffices to integrate the original, unreduced flow of Figure 10.1, but plot the solution in the image space, i.e., u_i invariant, Hilbert polynomial coordinates, as in figure 10.13. A drawback of such polynomial projections is that the folding mechanism is harder to view since the dynamics is squeezed near the z -axis.

Reducing dimensionality of a dynamical system by elimination of variables through inclusion of syzygies such as (10.50) introduces singularities. Such elimination of variables, however, is not needed for visualization purposes; syzygies merely guarantee that the dynamics takes place on a submanifold in the projection on the $\{u_1, u_2, \dots, u_m\}$ coordinates. However, when one reconstructs the dynamics in the original space M from its image M/G , the transformations have singularities at the fixed-point subspaces of the isotropy subgroups in M .

Nevertheless we can now easily identify a suitable Poincaré section, guided by the Lorenz flow examples of chapter 9, as one that contains the z -axis and the image of the relative equilibrium TW_1 , here defined by the condition $u_1 = u_4$. As in example 11.4, we construct the first return map using as coordinate the Euclidean arclength along the intersection of the unstable manifold of TW_1 with the Poincaré surface of section, see figure 10.14. Thus the goals set into the introduction to this chapter are attained: we have reduced the messy strange attractor of figure 10.1 to a 1-dimensional return map. As will be explained in




Figure 10.14: Return map to the Poincaré section $u_1 = u_4$ for complex Lorenz equations projected on invariant polynomials (10.49). The return map coordinate is the Euclidean arclength distance from TW_1 , measured along the Poincaré section of its spiral-out unstable manifold, as for the Lorenz flow return map of example 11.4.

example 11.4 for the Lorenz attractor, we now have the symbolic dynamics and can compute as many relative periodic orbits of the complex Lorenz flow as we wish, missing none.

What limits the utility of Hilbert basis methods are not such singularities, but rather the fact that the algebra needed to determine a Hilbert basis becomes computationally prohibitive as the dimension of the system or of the group increases. Moreover, even if such basis were available, rewriting the equations in an invariant polynomial basis seems impractical, so Hilbert basis computations appear not feasible beyond state space dimension of order ten. When our goal is to quotient continuous symmetries of high-dimensional flows, such as the Navier-Stokes flows, we need a more practical, workable framework. The method of moving frames of sect. 10.4 is one such minimalist alternative.

Résumé

The message: If a dynamical systems has a symmetry, use it! Here we have described how, and offered two approaches to continuous symmetry reduction. In the *method of slices* one fixes a ‘slice’ $(y - y')^T t' = 0$, a hyperplane normal to the group tangent t' that cuts across group orbits in the neighborhood of the slice-fixing point y' . Each class of symmetry-equivalent points is represented by a single point, with the symmetry-reduced dynamics in the reduced state space M/G given by (10.47):

$$u = v - \dot{\theta} \cdot t, \quad \dot{\theta} = (v \cdot t') / (t \cdot t').$$

In practice one runs the dynamics in the full state space, and post-processes the trajectory by the method of moving frames. In the *Hilbert polynomial basis* approach one transforms the equivariant state space coordinates into invariant ones, by a nonlinear coordinate transformation

$$\{x_1, x_2, \dots, x_d\} \rightarrow \{u_1, u_2, \dots, u_m\},$$

and study the invariant ‘image’ of dynamics (10.51) rewritten in terms of invariant coordinates.

In practice, continuous symmetry reduction is considerably more involved than the discrete symmetry reduction to a fundamental domain of chapter 9. Slices are only local sections of group orbits, and Hilbert polynomials are non-unique and difficult to compute for high-dimensional flows. However, there is no need to actually recast the dynamics in the new coordinates: either approach can be used as a visualization tool, with all computations carried out in the original coordinates, and then, when done, projecting the solutions onto the symmetry reduced state space by post-processing the data. The trick is to construct a good set of symmetry invariant Poincaré sections (see sect. 3.1), and that is always a dark art, with or without a symmetry.

In sect. 2.1.1 we made an attempt to classify ‘all possible motions:’ (1) equilibria, (2) periodic orbits, (3) everything else. Now one can discern in the fog of dynamics an outline of a more serious classification - long time dynamics takes place on the closure of a set of all invariant compact sets preserved by the dynamics, and those are: (1) 0-dimensional equilibria M_{EQ} , (2) 1-dimensional periodic orbits M_p , (3) global symmetry induced N -dimensional relative equilibria M_{TW} , (4) $(N+1)$ -dimensional relative periodic orbits M_p , (5) terra incognita. We have some inklings of the ‘terra incognita:’ for example, symplectic symmetry induces existence of KAM-tori, and in general dynamical settings we encounter *partially hyperbolic invariant tori*, isolated tori that are consequences of dynamics, not of a global symmetry. They are harder to compute than anything we have attempted so far, as they cannot be represented by a single relative periodic orbit, but require a numerical computation of full $(N+1)$ -dimensional compact invariant sets and their infinite-dimensional linearized Jacobian matrices, marginal in $(N+1)$ dimensions, and hyperbolic in the rest.

We conclude with a few general observations: Higher dimensional dynamics requires study of compact invariant sets of higher dimension than 0-dimensional equilibria and 1-dimensional periodic orbits studied so far. We expect, for example, partially hyperbolic invariant tori to play important role. In this chapter we have focused on the simplest example of such compact invariant sets, where invariant tori are a robust consequence of a global continuous symmetry of the dynamics. The direct product structure of a global symmetry that commutes with the flow enables us to reduce the dynamics to a desymmetrized $(d-1-N)$ -dimensional reduced state space M/G .

Relative equilibria and relative periodic orbits are the hallmark of systems with continuous symmetry. Amusingly, in this extension of ‘periodic orbit’ theory from unstable 1-dimensional closed periodic orbits to unstable $(N+1)$ -dimensional compact manifolds M_p invariant under continuous symmetries, there are either no or proportionally few periodic orbits. In presence of a continuous symmetry, likelihood of finding a periodic orbit is *zero*. Relative periodic orbits are almost never eventually periodic, i.e., they almost never lie on periodic trajectories in the full state space, so looking for periodic orbits in systems with continuous symmetries is a fool’s errand.

However, dynamical systems are often equivariant under a combination of continuous symmetries and discrete coordinate transformations of chapter 9, for example the orthogonal group $O(n)$. In presence of discrete symmetries relative periodic orbits within discrete symmetry-invariant subspaces are eventually periodic. Atypical as they are (no generic chaotic orbit can ever enter these discrete invariant subspaces) they will be important for periodic orbit theory, as there the shortest orbits dominate, and they tend to be the most symmetric solutions.

chapter 21

Commentary

Remark 10.1 A brief history of relativity, or, ‘Desymmetrization and its discontents’ (after Civilization and its discontents; continued from remark 9.1): The literature on symmetries in dynamical systems is immense, most of it deliriously unintelligible. Would it kill them to say ‘symmetry group of orbit \mathcal{M}_p ’ instead of carrying on about ‘isotropies, quotients, factors, normalizers, centralizers and stabilizers?’ For the dynamical systems applications at hand we need only basic results, on the level of any standard group theory textbook [10.2]. Chapter 2. of ref. [10.3] offers a pedagogical introduction to Lie groups of transformations, and Nakahara [10.4] to Lie derivatives and brackets. The presentation given here is in part based on Siminos thesis [10.29] and ref. [10.30]. The reader is referred to the monographs of Golubitsky and Stewart [10.42], Hoyle [10.44], Olver [10.45], Bredon [10.19], and Krupa [10.46] for more depth and rigor than would be wise to wade into here.

The relative equilibria and relative periodic solutions are related to equilibria and periodic solutions respectively of dynamics reduced by the symmetries. They appear in many physical situations, such as motion of rigid bodies, gravitational N -body problems, molecules, nonlinear waves, spiralling patterns and turbulence. According to Cushman, Bates [10.5] and Yoder [10.6], C. Huygens [10.7] understood the relative equilibria of a spherical pendulum many years before publishing them in 1673. A reduction of the translation symmetry was obtained by fixing the center of mass at the origin or, considering a configuration as an element of the disposition space $\mathbb{R}^D/(1, \dots, 1)\mathbb{R}$ by Jacobi. According to Chenciner [10.8, 10.10], the first attempt to find (relative) periodic solutions of the N -body problem by was the 1896 short note by Poincaré [10.9], in the context of the 3-body problem. Relative equilibria of the N -body problem (known in this context as the Lagrange points, stationary in the co-rotating frame) are circular motions in the inertial frame, and relative periodic orbits correspond to quasiperiodic motions in the inertial frame. Relative equilibria can exist in a rotating frame, and are called central configurations. For relative periodic orbits in celestial mechanics see also ref. [10.11]. Striking application of relative periodic orbits has been the discovery of “choreographies” of N -body problems [10.12, 10.13, 10.14].

The notion of a *moving frame* as a map from a manifold to a Lie group was introduced by Cartan [10.33], and the factorization (10.40) is stated (without reference) on p. 31 of Anosov and Arnol’d [10.34]. We follow Fels and Olver who view the method as an alternative to the Gröbner bases methods (here called ‘Hilbert bases’) to compute functionally independent fundamental invariant bases for general group actions, with no explicit connection to dynamics, differential equations or symmetry reduction. ‘Fundamental’ here means that they can be used to generate all other invariants. Olver’s monograph [10.45] is pedagogical, but does not describe the original Cartan’s method. Fels and Olver papers [10.40, 10.41] are lengthy and technical. They refer to Cartan’s method as method of ‘moving frames’ and view it as a special and less rigorous case of their ‘moving coframe’

method. The name ‘moving coframes’ arises through the use of Maurer-Cartan form which is a coframe on the Lie group G , i.e., they form a pointwise basis for the cotangent space. In refs. [10.29, 10.30] the invariant bases generated by the moving frame method are used as a basis to project a full state space trajectory to the slice (i.e., the \mathcal{M}/G reduced state space).

The basic idea of the ‘method of slices’ is intuitive; for example, it is stated without attribution as the problem 1. of Sect. 6.2 of Arnol’d *Ordinary Differential Equations* [10.32]. It is reinvented often,

HaMe98

Derivation of sect. 10.4.2 follows Rowley and Marsden [10.37] who, in the pattern recognition and ‘template fitting’ settings, call (10.46) the ‘reconstruction equation’ [10.36, 10.35].

For the definition of ‘slice’ see, for example, Chossat and Lauterbach [10.39]. Briefly, a submanifold \mathcal{M}_y containing y is called a *slice* through y if it is invariant under isotropy $G_y(\mathcal{M}_y) = \mathcal{M}_y$. If y is a fixed point of G , then slice is invariant under the whole group. The slice theorem is explained, for example, in Encyclopaedia of Mathematics. Slices tend to be discussed in contexts much more difficult than our application - symplectic groups, sections in absence of global charts, non-compact Lie groups. We follow refs. [10.35] in referring to a local group-orbit section as a ‘slice.’ Refs. [10.19, 10.16] and others refer to global group-orbit sections as ‘cross-sections,’ a term that we rather avoid, as it already has a different and well established meaning in physics. Duistermaat and Kolk [10.15] refer to ‘slices,’ but the usage goes back at least to Guillemin and Sternberg [10.16] in 1984, Palais [10.18] in 1961 and Mastow [10.17] in 1957. Bredon [10.19] discusses both cross-sections and slices. Guillemin and Sternberg [10.16] define the ‘cross-section,’ but emphasize that finding it is very rare: “existence of a global section is a very stringent condition on a group action. The notion of ‘slice’ is weaker but has a much broader range of existence.”

The Gilmore and Lettelier monograph [10.21] offers a very clear, detailed and user friendly discussion of symmetry reduction by means of Hilbert polynomial bases (do not look for ‘Hilbert’ in the index, though). The determination of a Hilbert basis appears computationally prohibitive for state space dimensions larger than ten [10.38, 10.39], and rewriting the equations of motions in invariant polynomial bases appears impractical for high-dimensional flows.

Sect. 10.1.2 ‘Lie groups for cyclists’ is bit of a joke in more ways than one. First, ‘cyclist,’ ‘pedestrian’ throughout ChaosBook.org refer jokingly both to the title of Lipkin’s *Lie groups for pedestrians* [10.22] and to our preoccupations with actual cycling. Lipkin’s ‘pedestrian’ is fluent in Quantum Field Theory, but wobbly on Dynkin diagrams. More to the point, it’s impossible to dispose of Lie groups in a page of text. As a counterdote to the 1-page summary of sect. 10.1.2, consider reading Gilmore’s monograph [10.20] which offers a quirky, personal and enjoyable distillation of a lifetime of pondering Lie groups. As seems to be the case with any textbook on Lie groups, it will not help you with the problem at hand, but it is the only place you can learn both what Galois actually did when he invented the theory of finite groups in 1830, and what, inspired by Galois, Lie actually did in his 1874 study of symmetries of ODEs. Gilmore also explains many things that we pass over in silence here, such as matrix groups, group manifolds, and compact groups.

Remark 10.2 Complex Lorenz equations (10.1) were introduced by Gibbon and McGuinness [10.24, 10.25] as a low-dimensional model of baroclinic instability in the at-

mosphere. They are a generalization of Lorenz equations (2.12). Ning and Haken [10.26] have shown that equations isomorphic to complex Lorenz equations also appear as a truncation of Maxwell-Bloch equations describing a single mode, detuned, ring laser. They set $e + \rho_2 = 0$ so that a detuned equilibrium exists. Bakasov and Abraham [10.27] criticize this choice as being “degenerate” (the choice of Ning and Haken leads to non-generic bifurcations) and show that one can use complex Lorenz equations with $\rho_2 = 0$ and $e \neq 0$ to describe detuned lasers. Here we are not interested in the physical applications of these equations; rather, we study them as a simple example of a dynamical system with continuous (but no discrete) symmetries, with a view of testing methods of reducing the dynamics to a lower-dimensional reduced state space. Complex Lorenz flow examples and exercises in this chapter are based on E. Siminos thesis [10.29] and R. Wilczak project report [10.31].

(E. Siminos and P. Cvitanović)

Exercises

10.1. **Visualizations of the 5-dimensional complex Lorenz flow:** Plot complex Lorenz flow projected on any three of the five $\{x_1, x_2, y_1, y_2, z\}$ axes. Experiment with different visualizations.

10.2. **An SO(2)-equivariant flow with two Fourier modes:** Complex Lorenz equations (10.1) of Gibbon and McGuinness [10.24] have a degenerate 4-dimensional subspace, with SO(2) acting only in its lowest non-trivial representation. Here is a possible model, still 5-dimensional, but with SO(2) acting in the two lowest representations. Such models arise as truncations of Fourier-basis representations of PDEs on periodic domains. In the complex form, the simplest such modification of complex Lorenz equations may be the “2-mode” system

$$\begin{aligned} \dot{x} &= -\sigma x + \sigma x^* y \\ \dot{y} &= (r - z)x^2 - ay \\ \dot{z} &= \frac{1}{2}(x^2 y^* + x^{*2} y) - bz, \end{aligned} \tag{10.53}$$

where $x, y, r = r_1 + i r_2, a = 1 + i e$ are complex and z, b, σ are real. Rewritten in terms of real variables $x = x_1 + i x_2, y = y_1 + i y_2$ this is a 5-dimensional first order ODE system

$$\begin{aligned} \dot{x}_1 &= -\sigma x_1 + \sigma y_1 \\ \dot{x}_2 &= -\sigma x_2 + \sigma y_2 \\ \dot{y}_1 &= (\rho_1 - z)x_1^2 - r_2 x_2 - y_1 - e y_2 \end{aligned}$$

10.3. **SO(2) rotations in a plane:** Show by exponentiation (10.7) that the SO(2) Lie algebra element \mathbf{T} generates rotation g in a plane,

$$g(\theta) = e^{\mathbf{T}\theta} = \cos \theta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \sin \theta \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}. \tag{10.55}$$

10.4. **Invariance under fractional rotations.** Argue that if the isotropy group of the velocity field $v(x)$ is the discrete subgroup C_m of SO(2) rotations about an axis (let’s say the ‘z-axis’),

$$C^{1/m} v(x) = v(C^{1/m} x) = v(x), \quad (C^{1/m})^m = e,$$

the only non-zero components of Fourier-transformed equations of motion are a_{jm} for $j = 1, 2, \dots$. Argue that the Fourier representation is then the quotient map of the dynamics, \mathcal{M}/C_m . (Hint: this sounds much fancier than what is - think first of how it applies to the Lorenz system and the 3-disk pinball.)

10.5. **U(1) equivariance of complex Lorenz equations for finite angles:** Show that the vector field in complex

$$\begin{aligned} \dot{y}_2 &= \\ \dot{z} &= -bz + x_1 y_1 + x_2 y_2. \end{aligned} \tag{10.54}$$

Verify (10.54) by substituting $x = x_1 + i x_2, y = y_1 + i y_2, r = r_1 + i r_2, a = 1 + i e$ into the complex 2-mode equations (10.53).

10.6. **SO(2) equivariance of complex Lorenz equations for finite angles:** Show that complex Lorenz equations (10.2) are equivariant under rotation for finite angles.

10.7. **Stability matrix of complex Lorenz flow:** Compute the stability matrix (10.26) for complex Lorenz equations (10.2).

10.8. **Rotational equivariance, infinitesimal angles.** Show that complex Lorenz equations are equivariant under infinitesimal SO(2) rotations.

10.9. **Discover the equivariance of a given flow:** Suppose you were given complex Lorenz equations, but nobody told you they are SO(2) equivariant. More generally, you might encounter a flow without realizing that it has a continuous symmetry - how would you discover it?

10.10. **Equilibria of complex Lorenz equations:** Find all equilibria of complex Lorenz equations. Hint: Equilibria come either in the fixed Fix (G) subspace, or on a group orbit.

10.11. **Equilibria of complex Lorenz equations:** In exercise 10.10 we found only one equilibrium of complex Lorenz equations. The Ning and Haken [10.26] version of complex Lorenz equations (a truncation of Maxwell-Bloch equations describing a single mode ring laser) sets $e + \rho_2 = 0$ so that a detuned equilibrium exists. Test your routines on 2 cases: (a) $e = 0, \rho_2 = 0$. As discussed by Siminos [10.29], reality of parameters a, ρ in (10.1) implies existence of a discrete C_2 symmetry. (b) $e + \rho_2 = 0, e \neq 0$. You might want to compare results with those of Ning and Haken.

10.12. **Complex Lorenz equations in polar coordinates.** Rewrite complex Lorenz equations from Cartesian to polar coordinates, using $(x_1, x_2, y_1, y_2, z) = (r_1 \cos \theta_1, r_1 \sin \theta_1, r_2 \cos \theta_2, r_2 \sin \theta_2, z)$,

Lorenz equations (10.1) is equivariant under the unitary group U(1) acting on $\mathbb{R}^5 \cong \mathbb{C}^2 \times \mathbb{R}$ by

$$g(\theta)(x, y, z) = (e^{i\theta} x, e^{i\theta} y, z), \quad \theta \in [0, 2\pi]. \tag{10.56}$$

(E. Siminos)

10.6. **SO(2) equivariance of complex Lorenz equations for finite angles:** Show that complex Lorenz equations (10.2) are equivariant under rotation for finite angles.

10.7. **Stability matrix of complex Lorenz flow:** Compute the stability matrix (10.26) for complex Lorenz equations (10.2).

10.8. **Rotational equivariance, infinitesimal angles.** Show that complex Lorenz equations are equivariant under infinitesimal SO(2) rotations.

10.9. **Discover the equivariance of a given flow:** Suppose you were given complex Lorenz equations, but nobody told you they are SO(2) equivariant. More generally, you might encounter a flow without realizing that it has a continuous symmetry - how would you discover it?



10.10. **Equilibria of complex Lorenz equations:** Find all equilibria of complex Lorenz equations. Hint: Equilibria come either in the fixed Fix (G) subspace, or on a group orbit.

10.11. **Equilibria of complex Lorenz equations:** In exercise 10.10 we found only one equilibrium of complex Lorenz equations. The Ning and Haken [10.26] version of complex Lorenz equations (a truncation of Maxwell-Bloch equations describing a single mode ring laser) sets $e + \rho_2 = 0$ so that a detuned equilibrium exists. Test your routines on 2 cases: (a) $e = 0, \rho_2 = 0$. As discussed by Siminos [10.29], reality of parameters a, ρ in (10.1) implies existence of a discrete C_2 symmetry. (b) $e + \rho_2 = 0, e \neq 0$. You might want to compare results with those of Ning and Haken.

10.12. **Complex Lorenz equations in polar coordinates.** Rewrite complex Lorenz equations from Cartesian to polar coordinates, using $(x_1, x_2, y_1, y_2, z) = (r_1 \cos \theta_1, r_1 \sin \theta_1, r_2 \cos \theta_2, r_2 \sin \theta_2, z)$,

$$(r_1 \cos \theta_1, r_1 \sin \theta_1, r_2 \cos \theta_2, r_2 \sin \theta_2, z), \tag{10.57}$$

where $r_1 \geq 0, r_2 \geq 0$. Show that in polar coordinates the equations take form

$$\begin{pmatrix} \dot{r}_1 \\ \dot{\theta}_1 \\ \dot{r}_2 \\ \dot{\theta}_2 \\ \dot{z} \end{pmatrix} = \begin{pmatrix} -\sigma(r_1 - r_2 \cos \theta) \\ -\sigma \frac{r_2}{r_1} \sin \theta \\ -r_2 + r_1((\rho_1 - z) \cos \theta - \rho_2 \sin \theta) \\ e + \frac{r_2}{r_1}((\rho_1 - z) \sin \theta + \rho_2 \cos \theta) \\ -bz + r_1 r_2 \cos \theta \end{pmatrix}, \tag{10.58}$$

10.13. **Visualizations of the complex Lorenz flow in polar coordinates:** Plot complex Lorenz flow projected on any three of the $\{r_1, r_2, \theta, z\}$ coordinates. Experiment with different visualizations. The flow (10.58) is singular as $r_j \rightarrow 0$, with angle θ_j going through a rapid change there: is that a problem? Does it make sense to insist on $r_1 \geq 0, r_2 \geq 0$, or should one let them have either sign in order that the θ trajectory be continuous?


10.14. **Computing the relative equilibrium TW₁:** Find the relative equilibria of the complex Lorenz equations by finding the equilibria of the system in polar coordinates (10.58). Show that

(a) The relative equilibrium (hereafter referred to [10.29] as TW₁) is given by

We know from classical mechanics that for translationally or rotationally invariant flows the relative distance is invariant (that is why one speaks of ‘relative’ equilibria), hence we introduce a variable $\theta = \theta_1 - \theta_2$. θ_1 and θ_2 change in time, but at the relative equilibria the difference between them is constant. Show that this new variable allows us to rewrite the complex Lorenz equations as 4 coupled polar coordinates equations:

$$\begin{pmatrix} \dot{r}_1 \\ \dot{r}_2 \\ \dot{\theta} \\ \dot{z} \end{pmatrix} = \begin{pmatrix} -\sigma(r_1 - r_2 \cos \theta) \\ -r_2 + (\rho_1 - z)r_1 \cos \theta \\ -e - \left(\sigma \frac{r_2}{r_1} + (\rho_1 - z) \frac{r_2}{r_1}\right) \sin \theta \\ -bz + r_1 r_2 \cos \theta \end{pmatrix} \tag{10.58}$$

where we have set $\rho_2 = 0$. Plot a long-time solution of these equations and show that the polar representation introduces singularities into what initially was a smooth flow:



We shall encounter the same problem in implementing the $x_1 = 0$ moving frames slice). A polar coordinates $\{r_1, r_2, \theta\}$ plot of the complex Lorenz flow strange attractor. θ is very small until the trajectory approaches either $r_1 \rightarrow 0$ or $r_2 \rightarrow 0$, where Mathematica continues through the singularity by a rapid change of θ by π . The the fixed Fix (G) subspace $(r_1, r_2, \theta, z) = (0, 0, \theta, z)$ separates the two folds of the attractor.

10.13. **Visualizations of the complex Lorenz flow in polar coordinates:** Plot complex Lorenz flow projected on any three of the $\{r_1, r_2, \theta, z\}$ coordinates. Experiment with different visualizations. The flow (10.58) is singular as $r_j \rightarrow 0$, with angle θ_j going through a rapid change there: is that a problem? Does it make sense to insist on $r_1 \geq 0, r_2 \geq 0$, or should one let them have either sign in order that the θ trajectory be continuous?

10.14. **Computing the relative equilibrium TW₁:** Find the relative equilibria of the complex Lorenz equations by finding the equilibria of the system in polar coordinates (10.58). Show that

(a) The relative equilibrium (hereafter referred to [10.29] as TW₁) is given by

$$(r_1, r_2, \theta, z) = \left(\sqrt{b(\rho_1 - d)}, \sqrt{bd(\rho_1 - d)}, \cos^{-1}\left(\frac{1}{\sqrt{d}}\right), \rho_1 - d \right), \quad (10.59)$$

where $d = 1 + e^2/(\sigma + 1)^2$,

(b) The angular velocity of relative equilibrium TW_1 is

$$\dot{\theta}_t = \sigma e/(\sigma + 1), \quad (10.60)$$

with the period $T_{TW_1} = 2\pi(\sigma + 1)/\sigma e$.

10.15. **Relative equilibrium TW_1 in polar coordinates:** Plot the equilibrium TW_1 in polar coordinates.

10.16. **Relative equilibrium TW_1 in Cartesian coordinates:** Show that for our parameter values,

$$\begin{aligned} x_{TW_1} &= (x_1, x_2, y_1, y_2, z) & (10.61) \\ &= (8.4849, 0.077135, 8.4856, 0, 26.999), \end{aligned}$$

is a point on the TW_1 orbit. Plot the relative equilibrium TW_1 in Cartesian coordinates.

10.17. **Eigenvalues and eigenvectors of TW_1 stability matrix:** Compute the eigenvalues and eigenvectors of the stability matrix (10.26) evaluated at TW_1 and using the (10.2) parameter values, in (a) Cartesian coordinates, (b) polar coordinates.

10.18. **The eigen-system of TW_1 stability matrix in polar coordinates:** Plot the eigenvectors of A at TW_1 in polar coordinates, as well as the complex Lorenz flow at values very near TW_1 .

10.19. **Eigenvalues and eigenvectors of EQ_0 stability matrix:** Find the eigenvalues and the eigenvectors of the stability matrix A (10.26) at $EQ_0 = (0, 0, 0, 0, 0)$ determined in exercise 10.10. ChaosBook convention is to order eigenvalues from most positive (unstable) to the most negative. Follow that. Replace complex eigenvectors by the real, imaginary parts, as that is what you actually use.

10.20. **The eigen-system of the stability matrix at EQ_0 :** Plot the eigenvectors of A at EQ_0 and the complex Lorenz flow at values very close to EQ_0 .

10.21. **SO(2) or harmonic oscillator slice:** Construct a moving frame slice for action of SO(2) on \mathbb{R}^2

$$(x, y) \mapsto (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$$

by, for instance, the positive y axis: $x = 0, y > 0$. Write out explicitly the group transformations that bring any point back to the slice. What invariant is preserved by this construction? (E. Siminos)

10.22. **State space reduction by a slice, finite time segments:** Replace integration of the complex Lorenz equations by a sequence of finite time steps, each followed by a rotation such that the next segment initial point is in the slice $y_2 = 0, y_1 > 0$.

10.23. **State space reduction by a slice, ODE formulation:** Reconsider (10.22) in the sequence of infinitesimal time steps limit, each followed by an infinitesimal rotation such that the next segment initial point is in the slice $y_2 = 0, y_1 > 0$. Derive the corresponding $4d$ reduced state space ODE for the complex Lorenz flow. Here is a way to do it, bit different from the derivation given in sect. 10.4.2.


Infinitesimal time version of the moving frames symmetry reduction is attained by taking small time steps in figure 10.11 and dropping the higher order terms. For infinitesimal $d\theta$ we set $\sin d\theta \approx d\theta, \cos d\theta \approx 1, g(d\theta) \approx 1 + d\theta \mathbf{T}$, and the condition (10.41) for rotating an infinitesimal time evolution step $dx = v dt$ back into the slice

$$\begin{aligned} 0 &= (y + dx) \cdot g(d\theta)^T \mathbf{T} y' \\ &\approx (y + dt v) \cdot (1 + d\theta \mathbf{T})^T \mathbf{T} y' \\ &\approx dt v \cdot \mathbf{T} y' + d\theta y \cdot \mathbf{T}^T \mathbf{T} y' \end{aligned}$$

yields

$$d\theta \approx -\frac{v \cdot \mathbf{T} y'}{y \cdot \mathbf{T}^T \mathbf{T} y'} dt. \quad (10.62)$$

Let $u(y)$ be the vector field that generates the flow in the reduced state space. According to



in the limit that $g(d\theta) \approx 1 + d\theta \mathbf{T}$ the infinitesimal time step under u is connected to the time step under v by

$$y + u dt = (1 + d\theta \mathbf{T}) \cdot (y + v dt).$$

Dropping second order terms, dividing through with dt

$$u = v + \frac{d\theta}{dt} \mathbf{T} y,$$

and substituting (10.62) gives the reduced state space equations (10.47):

$$\dot{x} = v - \frac{(v \cdot \mathbf{T} y')}{(y \cdot y')_4} \mathbf{T} y, \quad (10.63)$$

where we have used the fact that $-x \cdot \mathbf{T} \mathbf{T} x^* = (x \cdot x^*)_4 = x_1 x_1^* + x_2 x_2^* + y_1 y_1^* + y_2 y_2^*$ is the dot-product restricted to the 4-dimensional representation of SO(2). By construction, the motion stays in the $(d-1)$ -dimensional slice.

10.24. **Accumulated phase shift:** Derive the $1d$ equation (10.46) for the accumulated phase shift θ associated with the 4-dimensional reduced state space ODE of exercise 10.23.

10.25. **The moving frame flow stays in the reduced state space:** Show that the flow (10.63) stays in a $(d-1)$ -dimensional slice.

10.26. **State space reduction by a relative equilibrium TW_1 cross-section:** Replace integration of the complex Lorenz equations by a sequence of short time steps, each followed by a rotation such that the next segment initial point is in the relative equilibrium TW_1 cross-section

$$(y - y_{TW_1}) \cdot t_{TW_1} = 0, \quad t_{TW_1} = \mathbf{T} y_{TW_1}, \quad (10.64)$$

where for any $x, y = g(\theta) \cdot x$ is the rotation that lies in the cross-section. Check figure 10.12 by long-time integration of (10.63).

References

[10.1] P. Cvitanović, R. L. Davidchack and E. Siminos, “State space geometry of a spatio-temporally chaotic Kuramoto-Sivashinsky flow,” *SIAM J. Applied Dynam. Systems* (2009); arXiv:0709.2944.

[10.2] M. Hamermesh, *Group theory and its application to physical problems* (Addison-Wesley, Reading, 1962).

[10.3] G. W. Bluman and S. Kumei, *Symmetries and differential equations* (Springer, New York 1989).

[10.4] M. Nakahara, *Geometry, topology and physics* (Inst. of Physics Publ., Bristol, 1990).

[10.5] R. H. Cushman and L. M. Bates, *Global aspects of classical integrable systems*, p. 402 (Birkhäuser, Basel 1997).

[10.6] J. G. Yoder, *Unrolling time: Christiaan Huygens and the mathematization of nature* (Cambridge Univ. Press, Cambridge 1988).

[10.7] C. Huygens, *L’Horloge à pendule* (Swets & Zeitlinger, Amsterdam 1673, 1967).

[10.8] A. Albouy and A. Chenciner, “Le problème des n corps et les distances mutuelles,” *Invent. Math.* **31**, 151 (1998).

[10.9] H. Poincaré, “Sur les solutions périodiques et le principe de moindre action,” *C.R.A.S.* **123**, 915 (1896).

[10.10] A. Chenciner, “Symmetries and ‘simple’ solutions of the classical N -body problem,” in *Proceedings ICMP03* (2003).

[10.11] R. Broucke, “On relative periodic solutions of the planar general three-body problem,” *Celestial Mechanics and Dynamical Astronomy* **12**, 439 (1975).

[10.12] A. Chenciner and R. Montgomery, “A remarkable solution of the 3-body problem in the case of equal masses,” *Ann. Math.* **152**, 881 (2000).

- [10.13] A. Chenciner, J. Gerver, R. Montgomery and C. Simó, "Simple choreographic motions of N -bodies: A preliminary study," pp. 287-308 in P. Newton, P. Holmes and A. Weinstein, eds., *Geometry, mechanics and dynamics* (Springer, New York 2002).
- [10.14] C. McCord, J. Montaldi, M. Roberts and L. Sbano, "Relative periodic orbits of symmetric Lagrangian systems," pp. 482-493, in Freddy Dumortier and et.al., eds., *Proceedings of Equadiff 2003* (2004).
- [10.15] J. J. Duistermaat and J. A. C. Kolk, *Lie groups* (Springer, New York 2000).
- [10.16] V. Guillemin and S. Sternberg, *Symplectic techniques in physics* (Cambridge Univ. Press, Cambridge 1990).
- [10.17] G. D. Mostow, "Equivariant embeddings in Euclidean space," *Ann. Math.* **65**, 432 (1957).
- [10.18] R. S. Palais, "On the existence of slices for actions of non-compact Lie groups," *Ann. Math.* **73**, 295 (1961).
- [10.19] G. Bredon, *Introduction to compact transformation groups*, (Academic Press, New York 1972).
- [10.20] R. Gilmore, *Lie groups, physics, and geometry* (Cambridge Univ. Press, Cambridge 2008).
- [10.21] R. Gilmore and C. Letellier, *The symmetry of chaos* (Oxford U. Press, Oxford 2007).
- [10.22] H. J. Lipkin, *Lie groups for pedestrians* (North-Holland, Amsterdam 1966).
- [10.23] T. Hawkins, *Emergence of the theory of Lie groups: An essay in the history of mathematics, 1869-1926* (Springer, New York 2000).
- [10.24] J. Gibbon and M. McGuinness, "The real and complex Lorenz equations in rotating fluids and lasers," *Physica D* **5**, 108 (1982).
- [10.25] A. Fowler, J. Gibbon and M. McGuinness, "The complex Lorenz equations," *Physica D* **4**, 139 (1982).
- [10.26] C.-z. Ning and H. Haken, "Detuned lasers and the complex Lorenz equations: Subcritical and supercritical Hopf bifurcations," *Phys. Rev.* **41**, 3826 (1990).
- [10.27] A. A. Bakasov and N. B. Abraham, "Laser second threshold: Its exact analytical dependence on detuning and relaxation rates," *Phys. Rev. A* **48**, 1633 (1993).
- [10.28] N. B. Abraham, U. A. Allen, E. Peterson, A. Vicens, R. Vilaseca, V. Espinosa and G. L. Lippi, "Structural similarities and differences among attractors and their intensity maps in the laser-Lorenz model," *Optics Comm.* **117**, 367 (1995).

- [10.29] E. Siminos, *Recurrent spatio-temporal structures in presence of continuous symmetries*, Ph.D. thesis (Georgia Inst. of Tech. 2009); ChaosBook.org/projects/theses.html.
- [10.30] E. Siminos and P. Cvitanović, "Continuous symmetry reduction and return maps for high dimensional flows," *Physica D*, (2010).
- [10.31] R. Wilczak, *Reducing the state-space of the complex Lorenz flow*, NSF REU summer 2009 project, U. of Chicago (2009); ChaosBook.org/projects/Wilczak/blog.pdf.
- [10.32] V.I. Arnol'd, *Ordinary differential equations* (Springer-Verlag, New York 1992).
- [10.33] É. Cartan, "La méthode du repère mobile, la théorie des groupes continus, et les espaces généralisés," *Exposés de Géométrie* **5** (1935).
- [10.34] D. V. Anosov and V. I. Arnol'd, *Dynamical systems I: Ordinary Differential Equations and Smooth Dynamical Systems* (Springer, 1988).
- [10.35] C. W. Rowley and J. E. Marsden, "Reconstruction equations and the Karhunen-Loève expansion for systems with symmetry," *Physica D* **142**, 1 (2000).
- [10.36] J. E. Marsden and T. S. Ratiu, *Introduction to mechanics and symmetry* (Springer, New York, 1994).
- [10.37] C. W. Rowley, I. G. Kevrekidis, J. E. Marsden and K. Lust, "Reduction and reconstruction for self-similar dynamical systems," *Nonlinearity* **16**, 1257 (2003).
- [10.38] K. Gatermann, *Computer algebra methods for equivariant dynamical systems* (Springer-Verlag, New York 2000).
- [10.39] P. Chossat and R. Lauterbach, *Methods in equivariant bifurcations and dynamical systems* (World Scientific, Singapore 2000).
- [10.40] M. Fels and P. J. Olver, "Moving coframes: I. a practical algorithm," *Acta Appl. Math.* **51**, 161 (1998).
- [10.41] M. Fels and P. J. Olver, "Moving coframes: II. regularization and theoretical foundations," *Acta Appl. Math.* **55**, 127 (1999).
- [10.42] M. Golubitsky and I. Stewart, *The symmetry perspective* (Birkhäuser, Boston 2002).
- [10.43] M. Golubitsky, I. Stewart and D. G. Schaeffer, *Singularities and groups in bifurcation theory, vol. II* (Springer-Verlag, New York 1988).
- [10.44] R. Hoyle, *Pattern formation: An introduction to methods* (Cambridge Univ. Press, Cambridge 2006).
- [10.45] P. J. Olver, *Classical invariant theory* (Cambridge Univ. Press, Cambridge 1999).

- [10.46] M. Krupa, "Bifurcations of relative equilibria," *SIAM J. Math. Anal.* **21**, 1453 (1990).
- [10.47] D. Viswanath, "Recurrent motions within plane Couette turbulence," *J. Fluid Mech.* **580**, 339 (2007); [arXiv:physics/0604062](https://arxiv.org/abs/physics/0604062).